

ON THE COMPLEXITY OF H -COLOURING PLANAR GRAPHS

G.MACGILLIVRAY AND M.SIGGERS

ABSTRACT. We show that if H is an odd-cycle, or any non-bipartite graph of girth 5 and maximum degree at most 3, then planar H -COL is NP -complete

1. INTRODUCTION

Given a graph H , the problem of whether or not another graph G admits an H -colouring, i.e., an edge-preserving vertex map to H , is referred to as H -COL. In [10], it was shown that H -COL is NP -complete if H contains an odd cycle, and polynomial-time solvable otherwise. (In this note, all graphs are assumed to be finite, symmetric, and irreflexive.)

One can ask whether the H -COL problem remains NP -complete when the instances G are given certain restrictions. A good reference for such problems is [9].

In this paper we focus on restricting to planar graphs G . We refer to the problem of H -COL, restricted to planar instances, as *planar H -COL*. In [5], it is shown that planar K_3 -COL is NP -complete. At the same time, the four-colour theorem implies that planar H -COL is trivial for any graph H containing a K_4 .

In general, it seems to be a difficult problem to determine for which H , planar H -COL is NP -complete. On the one hand, it has recently been shown in [8] that planar H -COL is NP -complete for any non-bipartite planar graph not containing a K_4 . On the other hand, in [12], it is shown that a planar graph G maps to the Clebsch graph C if and only if it is K_3 -free. This implies that planar C -COL is polynomial time solvable. This result has recently been extended in [13] to show that, in particular, for any planar graph F there is a graph U containing no homomorphic image of F , for which there is a U -colouring for any F -free graph G . Planar U -COL is polynomial time solvable for any such graph U .

Restriction to planar instances has been considered in other homomorphism problems. In [11], the complexity of the list-colouring problem is investigated for planar instances with certain other restrictions, such as list size bounds and degree bounds. In [3] the problem of acyclic digraph homomorphisms was investigated for planar instances.

In this note, we prove the following two theorems.

Theorem 1.1. *For any odd integer $\ell \geq 5$, planar C_ℓ -COL is NP -complete.*

Theorem 1.2. *Let H be any graph of girth 5 and maximum degree 3, then planar H -COL is NP -complete.*

Theorem 1.1 has recently been proved independently in [8], using a different proof. In the same paper, as is mentioned above, they also characterise for which

Date: August 29, 2006.

planar graphs H the problem of planar H -COL is NP -complete. Our Theorem 1.2 is of interest, as it shows that planar H -COL is NP -complete for many non-planar graphs H . For example, it implies that planar H -COL is NP -complete when H is the Petersen graph.

Outline of the paper. In Section 2 we prove Theorem 1.1. In the other three sections we prove Theorem 1.2. In Section 3 we provide some reductions which will allow us to assume certain useful properties about the graphs H of Theorem 1.2. In Section 4 we use the assumptions about H to provide a structural description of such graphs. In Section 5, we prove that planar H -COL is NP -complete for any graph H satisfying this structural description.

Notation. Throughout the paper, when we consider a C_ℓ -colouring of a graph, we will assume that the vertices of C_ℓ are $[\ell]$, the integers modulo ℓ , and that consecutive integers are adjacent. The same will be true for K_3 -colourings. For any set S , an S -colouring is a homomorphism to the complete graph having S as its vertex set. For any integer ℓ , an ℓ -path is a path consisting of ℓ vertices. It will often be necessary to consider a fixed embedding of a planar graph G , in this case we will refer to G as a *plane* graph. It is important to note that given a planar graph, a plane embedding can be found in polynomial time. See, for example, [1].

2. PROOF OF THEOREM 1.1

We prove Theorem 1.1 with an indicator-type construction, similar to the constructions of [10]. Given a planar graph G we will construct a planar graph $*G$, and show that

$$*G \rightarrow C_\ell \iff G \rightarrow K_3.$$

Because planar K_3 -COL is NP -complete, this implies the theorem.

To simplify notation, we assume G has no pendant edges. Clearly, removing pendant edges has no effect on whether or not G has a K_3 -colouring.

Before we get to Construction 2.3, which will give us $*G$, we define two graphs that we use in this, and other, constructions.

Definition 2.1. (The graphs $I(\ell)$ and $J(\ell)$).

Given an odd integer $\ell \geq 5$, define the graphs $I(\ell)$ and $J(\ell)$ as follows.



- The graph $I(\ell)$ is an $(\ell + 2)$ -cycle containing the path xv_1av_2y , plus the new path $xv'_1a'v'_2y$, introduced between the vertices x and y .
- The graph $J(\ell)$ is an ℓ -cycle containing the path xy , plus a new vertex b' which is joined to x and y .

The graphs $I(5)$ and $J(5)$ are pictured in Figure 1.

We will often use $I(\ell)$ and $J(\ell)$ in constructions in the following way.

Definition 2.2. Given a graph with vertices u and v we take a copy of $I = I(\ell)$ or $J = J(\ell)$ and identify u and v with the copies of x and y , respectively. When we do this, we will say that we **connect the vertices u and v with a copy of I (or J)**.

Observe the following properties of $I = I(\ell)$ and $J = J(\ell)$.

FIGURE 1. The graphs $I(5)$ and $J(5)$.FIGURE 2. Steps of Construction 2.3, where $G = K_3$ and $\ell = 5$.

P_I : A mapping f of $\{a, a', x, y\} \subset V(I)$ to $[\ell]$, can be extended to a C_ℓ -colouring of I if and only if $f(x)$ and $f(y)$ are distinct elements of the set $\{f(a) - 2, f(a), f(a) + 2\} \cap \{f(a') - 2, f(a'), f(a') + 2\}$.

P_J : J is C_ℓ colourable, and every C_ℓ -colouring assigns b and b' the same colour.

Admittedly, P_I is not the most natural characterisation of the colourings of I , but it will prove useful. We now give our indicator-type construction.

Construction 2.3. Given an odd integer $\ell \geq 5$ and a plane graph G without pendant edges, construct the plane graph $*G$ from G as follows.

- For every edge $e = uv$ of G , remove e and connect u and v with a copy I_e of $I = I(\ell)$. Do so with the embedding of I shown in Figure 1, so that the copies of the vertices a and a' are in different faces of the new graph.
- For every face F (including the outer face) of the original graph G , do the following. Let $e_1, e_2, \dots, e_{|F|}$ denote the edges of the boundary of F , ordered so that consecutive edges share a vertex. Let $a_{F,i}$ be the copy of a or a' in I_{e_i} that lies on the face F . For $i = 1, 2, \dots, |F| - 1$, connect the vertices $a_{F,i}$ and $a_{F,i+1}$ with a copy $J_{F,i}$ of $J = J(\ell)$.

See Figure 2 for an example of this construction. The figure shows the two steps of the construction, as applied to the $G = K_3$ with $\ell = 5$. The thick edges in the figure represent copies of J , with the endpoints of the thick edges representing the copies of the vertices b and b' .

It is clear that Construction 2.3 can be done so that the resulting graph $*G$ is planar. Furthermore, we use properties P_I and P_J to prove the following.

Lemma 2.4. *For any odd integer $\ell \geq 5$ and any plane graph G without pendant edges, let $*G$ be the planar graph returned by Construction 2.3. Let f be a map from $V(G)$ to $[\ell]$. Then f can be extended to a C_ℓ -colouring of $*G$ if and only if for every face F of G there exists $c_F \in [\ell]$ such that f induces a proper $\{c_F - 2, c_F, c_F + 2\}$ -colouring of the boundary of F .*

Proof. Assume that f can be extended to a C_ℓ -colouring, ϕ' , of $*G$. For each face F of G , the vertices $a_{F,i}$ are connected by copies of J , thus property P_J implies that $\phi'(a_{F,1}) = \cdots = \phi'(a_{F,|F|})$. Setting $c_F = \phi'(a_{F,1})$, property P_I implies us that ϕ' induces a $\{c_F - 2, c_F, c_F + 2\}$ -colouring of the boundary of F . Thus the same is true of f . This was for an arbitrary face F of G , so we have the forward implication of the lemma.

For the converse implication, let $f : V(G) \rightarrow [\ell]$, and let $F \mapsto c_F$ be a mapping of the faces of G to $[\ell]$, such that for every F , f induces a $\{c_F - 2, c_F, c_F + 2\}$ -colouring of the boundary of F . We extend f to a C_ℓ -colouring of $*G$ as follows.

For every edge $e = uv$ in G , let F_e and F'_e be the faces of G whose boundaries contain e . Assume, without loss of generality, that the copy of a in I_e lies in F_e , and the copy of a' lies in F'_e . Since u and v are adjacent in F , the definition of the map $F \mapsto c_F$ implies that, $f(u)$ and $f(v)$ are distinct members of both $\{c_{F_e} - 2, c_{F_e}, c_{F_e} + 2\}$ and $\{c_{F'_e} - 2, c_{F'_e}, c_{F'_e} + 2\}$. Thus by property P_I , f can be extended to C_ℓ -colouring of I_e in which $f(a) = c_{F_e}$ and $f(a') = c_{F'_e}$.

Now for any face F of G , we have that $f(a_{F,1}) = \cdots = f(a_{F,|F|}) = c_F$, thus by property P_J , f can be extended to a C_ℓ -colouring of $J_{F,i}$ for $i = 1, \dots, |F|$. We have thus C_ℓ -coloured $*G$, and so completed the proof of the lemma. \square

Now if ϕ is a K_3 -colouring of G , then defining $f : V(G) \rightarrow [\ell]$ by

$$f(v) = \begin{cases} 1 & \text{if } \phi(v) = 1 \\ 3 & \text{if } \phi(v) = 2 \\ \ell - 1 & \text{if } \phi(v) = 3 \end{cases},$$

Lemma 2.4 implies that there is a C_ℓ -colouring of $*G$ in which $f(a) = f(a') = 1$ for all copies of I . Thus $G \rightarrow K_3 \Rightarrow *G \rightarrow C_\ell$. We finish the proof of the theorem by proving the following lemma.

Lemma 2.5. *For any odd integer $\ell \geq 5$ and plane graph G , the planar graph $*G$ returned by Construction 2.3 has the property that*

$$*G \rightarrow C_\ell \Rightarrow G \rightarrow K_3.$$

Proof. Towards contradiction, assume that there is some plane graph G for which $*G \rightarrow C_\ell$, but $G \not\rightarrow K_3$. Further assume that G is a minimum such counter-example with respect to the number of vertices. Let ϕ' be a C_ℓ -colouring of $*G$.

By Lemma 2.4, ϕ' restricts to a 3-colouring (in G) of the boundary of any face of G . We may assume the following.

Claim 2.6. *Every face of G is a triangle.*

Proof. If some face is not a triangle, then $\phi'(u) = \phi'(v)$ for some vertices u and v in the boundary of the face. Let G' be the graph constructed from G by identifying u and v . (Remove any multiple edges thus introduced.) Clearly $G' \not\rightarrow K_3$.

We now show that there exists a C_ℓ -colouring of $*G'$. Thus G' will be a smaller counter-example to the lemma than G , which will contradict our choice of G . This will prove the claim.

By Lemma 2.4, ϕ' restricts to a map $f : V(G) \rightarrow [\ell]$, and induces a map $F \mapsto c_F \in [\ell]$ of the faces of G such that for any face F , f induces a proper $\{c_F - 2, c_F, c_F + 2\}$ -colouring of the boundary of F . Since $f(u) = f(v)$, f is a well defined map of $V(G')$. For any face F in G' , either the boundary of F is exactly the boundary of a face of G , or it is a path in the boundary of some face of G , in which the endpoints received the same colour under f , and have been identified. In either case, it is clear that f induces on it a $\{c - 2, c, c + 2\}$ -colouring for some c in $[\ell]$. Thus by Lemma 2.4, G' has a C_ℓ -colouring.

This proves the claim. \square

Now let x be a vertex of G . Since G is a triangulation, the neighbourhood of x is a cycle. For every edge uv in this cycle, Lemma 2.4 implies that $\{\phi'(u), \phi'(v)\}$ is equal to one of $\{\phi'(x) - 4, \phi'(x) - 2\}$, $\{\phi'(x) - 2, \phi'(x) + 2\}$, and $\{\phi'(x) + 2, \phi'(x) + 4\}$. That is to say, the neighbourhood of x in G is a cycle, and admits a homomorphism to the path $\phi'(x) - 4, \phi'(x) - 2, \phi'(x) + 2, \phi'(x) + 4$. Thus the neighbourhood of x in G is an even cycle, and so x has even degree. Since x was an arbitrary vertex of G , the fact that G is 3-colourable follows from the fact that any triangulation in which every vertex has even degree (i.e., any eulerian triangulation) is 3-colourable. This result is apparently from [7], and is given as an exercise (exer. 9.6.2) in [1].

The graph G being 3-colourable contradicts our original assumption, so this completes the proof of the lemma and of the theorem. \square

3. REDUCTIONS

Our goal in the remainder of the paper is to show that planar H -COL is NP -complete any for graph H of girth 5 and maximum degree 3. Recall that the *core* of a graph is the unique (up to isomorphism) maximal subgraph whose only monomorphisms are automorphisms. It is well known (see, for example, [9]) that H -COL is polynomially equivalent to C -COL, where C is the core of H .

Thus, in the proof of Theorem 1.2, we will consider only graphs H with the following properties.

- P1) H is a core.
- P2) H has girth 5.
- P3) The maximum degree of H is 3.

In this section we show that we may further assume that H has the following three properties.

- P4) H is connected.
- P5) Every edge of H is in a C_5 .
- P6) No vertex v of H , of degree 3, has every pair of neighbours in a C_5 .

We justify property P4 with the following lemma.

Lemma 3.1. *If planar H' -COL is NP -complete for every component H' of a graph H , then planar H -COL is NP -complete.*

Proof. The lemma is true if H has one component. Assume that the lemma is true for any graph with $d - 1$ components, and let H be a graph with d components

H_i such that planar H_i -COL is NP -complete for each of them. We show that the lemma is true for H , and so prove the lemma by induction.

Let H_1 be a component of H . Either every planar graph that admits an H_1 -colouring also admits an $(H - H_1)$ -colouring, or there is some planar graph P that admits an H_1 -colouring, but no $(H - H_1)$ -colouring. In the first case, planar $(H - H_1)$ -COL being NP -complete (by induction) implies that planar H -COL is.

For the second case, we provide a construction which reduces planar H_1 -COL to planar H -COL in polynomial time. This will imply that planar H -COL is NP -complete, so complete the proof of the lemma.

Since planar H_1 -COL is NP -complete, so is H_1 -COL, so by [10], H_1 contains an odd cycle. Thus there is some k such that for every pair of vertices in H_1 there is a walk between them of length k . Give an planar graph G , construct the planar graph G' as follows. For every vertex v of G , let P_v be a copy of P , and connect P_v to G with a path of length k .

Any H -colouring of G' induces an H_1 -colouring of G . Indeed, every copy of P must map to H_1 , so every vertex of the subgraph G of G' must also map to H_1 . On the other hand, one can check that any H_1 -colouring of G can be extended to an H -colouring of G' . This reduces planar H_1 -COL to planar H -COL, so planar H -COL is NP -complete. \square

To justify property P5, construct G' from given plane graph G as follows. For each edge of G , add a path of length four (involving three new vertices) joining its endpoints. The resulting graph, G' , is planar, and has the property that every edge belongs to a 5-cycle. Hence any homomorphism of G' to H must map its subgraph G to the subgraph H' of H induced by the edges of H belonging to 5-cycles. It follows that H' -colouring can be polynomially transformed to H -colouring. (This construction can be seen as a special case of the edge-sub-indicator construction from [10] (also see [9]). Hence, we have proved:

Lemma 3.2. *Let H be a graph of odd girth 5, and H' be its subgraph consisting of all edges of that are part of a C_5 in H . Then if planar H' -COL is NP -complete, so is planar H -COL.*

To justify property P6, we use a simple variation of Construction 2.3.

Lemma 3.3. *Let H be a graph of girth 5 and maximum degree 3. If H has the following property,*

There exists a vertex v of degree 3 in H such that for every pair w, w' of neighbours of v there is a C_5 in H containing the path $vw w'$.

then planar H -COL is NP -complete.


Before we prove the lemma, we present a definition used in its proof.

Definition 3.4. *For every vertex v in a graph H let \mathcal{N}_v be the graph defined by $V(\mathcal{N}_v) = N_H(v)$, and $E(\mathcal{N}_v) = \{uw \mid uvw \text{ is in a } C_5 \text{ in } H\}$.*

Proof of Lemma 3.3. Given any (non-empty) plane graph G , construct $*G$ exactly as in Construction 2.3 for $\ell = 5$, except use copies of $J(5)$ in place of copies of $I(5)$.

Claim 3.5. *For any graph H of odd girth 5,*

$$*G \rightarrow H \iff \exists v \in V(H) \text{ such that } G \rightarrow \mathcal{N}_v.$$

FIGURE 3. A sub-divided n -ribbon, S_n .

Proof. Let ϕ be an \mathcal{N}_v -colouring of G for some $v \in V(H)$. Define $\phi' : V(*G) \rightarrow V(H)$ as follows. If u is any copy of b or b' in any copy of $J(5)$ in $*G$, then let $\phi'(u) = v$. If $u \in V(G) \subset V(*G)$, then let $\phi'(u) = \phi(u)$. One can check that the map ϕ' can be extended to an H -colouring of $*G$.

On the other hand, let ϕ' be an H -colouring of $*G$. By property P_J , all copies of b and b' in $*G$ are mapped to the same vertex by ϕ' . Furthermore, for any edge uv in G , the vertices u and v are neighbours of some copy of b or b' in a copy of C_5 in $*G$. Thus they must map to distinct neighbours of $\phi'(b)$ in H that are adjacent in $\mathcal{N}_{\phi'(b)}$. That is to say, ϕ' restricted to $V(G)$ is an $\mathcal{N}_{\phi'(b)}$ -colouring of G . \square

The lemma follows. Indeed, since H has maximum degree 3, \mathcal{N}_v is a subgraph of K_3 for every $v \in V(H)$. If H has the property of the lemma, then $\mathcal{N}_v = K_3$ for some v . This allows us to rephrase the biconditional of the claim as

$$*G \rightarrow H \iff G \rightarrow K_3.$$

Since $*G$ is planar if G is, and planar K_3 -COL is NP -complete, this gives the conclusion of the lemma. \square

4. STRUCTURAL DESCRIPTION OF H

In this section, we give, with Lemma 4.6, a structural description of the graph H which have the properties P1-P6 listed at the beginning of Section 3. We begin with several definitions which we will need for the statement of Lemma 4.6.

Definition 4.1. (*Ribbons R_n and R_n^t .*) For $n \geq 3$, the **n -ribbon**, R_n , is the graph with vertices a_i and b_i for $i = 1, \dots, n$, and with edges $a_i b_i$, $a_i a_{i+1}$, and $b_i b_{i+1}$, (where indices are mod n) for $i = 1, \dots, n$. The **twisted n -ribbon**, R_n^t , is R_n with the edges $a_n b_1$ and $b_n a_1$ instead of $a_n a_1$ and $b_n b_1$.

The subgraph L_i induced by a_i, a_{i+1}, b_i , and b_{i+1} , is the i^{th} **link** of R_n or R_n^t .

Definition 4.2. (*Subdivided Ribbons S_n and S_n^t .*) A **subdivided n -ribbon**, S_n , is any subdivision of R_n in which exactly one edge of every link is subdivided into a 3-path. (See Figure 3.) S_n^t is an analagous subdivision of R_n^t . We further require that S_n and S_n^t have girth 5, (but this is only a restriction when $n = 3$ or 4).

Every link L_i of S_n or S_n^t thus contains 5 vertices. The fifth vertex, c_i is called the **codicil** of L_i . Such vertices may get more than one label, indeed if the codicil c_i of L_i is introduced in between a_{i+1} and b_{i+1} then it is the codicil of L_{i+1} as well, and so will also be referred to as c_{i+1} .

Observe that many distinct subdivided ribbons are isomorphic. That is, by reindexing the links of a graph S_n or by interchanging the sets $\{a_1, \dots, a_n\}$ and

$\{b_1, \dots, b_n\}$, we can get back a different subdivided n -ribbon. There are similar relabellings of S_n^t that allow us to make any link the ‘twisted’ link. This will often allow us to make certain assumptions about a graph S_n or S_n^t .

Definition 4.3. (*Link Types*) The link L_i of a (twisted) subdivided n -ribbon S_n or S_n^t is of type *LEFT*, *TOP*, *BOT*, or *RIGHT*, if c_i subdivides $a_i b_i$, $a_i a_{i+1}$, $a_{i+1} b_{i+1}$, or $b_i b_{i+1}$, respectively. In the twisted link L_n of S_n^t , types *TOP* and *BOT* must be redefined. The link is of type *TOP* if c_n subdivides $a_n b_1$, and of type *BOT* if c_n subdivides $b_n a_1$.

Definition 4.4. (*Broken Ribbons*) A **broken n -ribbon**, B_n , is a subgraph of a subdivided n -ribbon S_n with the edges, or 3-paths, between a_n and a_1 , and between b_n and b_1 , removed.

Observe that a broken n -ribbon has only $n - 1$ links, and up to relabelling, its first and last links can be assumed to be of type *LEFT* or *RIGHT*.

Definition 4.5. (*Caps C_A and C_B , and Bridges*) The **A-cap** C_A of a subdivided n -ribbon S_n is the cycle consisting of the vertices a_i , and all codicils of links of type *TOP*. The **B-cap** C_B of a subdivided n -ribbon S_n is the cycle consisting of the vertices b_i , and all codicils of links of type *BOT*.

The **A-cap** C_A of a twisted subdivided n -ribbon S_n^t is the cycle consisting of the vertices a_i , all codicils of links of type *TOP*, b_n , and c_n if L_n is of type *BOT*. The **B-cap** is defined similarly.

A **bridge** of S_n or S_n^t is a path $c_i x_i c_{i+1}$ with a new vertex x_i , where L_i is of type *TOP* and L_{i+1} is of type *BOT*, or vice versa. (When H contains a twisted subdivided n -ribbon S_n^t then we apply this definition to the n^{th} link by moving the twist to some non-adjacent link.)

We can finally give the structural description of a graph H satisfying properties P1 - P6. The rest of the section will be dedicated to proving this description.


Lemma 4.6. *Let H be a graph satisfying properties P1-P6. Then H contains a subdivided n -ribbon $S = S_n$ or S_n^t for some $n \geq 4$, with caps of girth at least 6. Its only other vertices or edges are in bridges of S .*

Proof. We begin by establishing the following claim.

Claim 4.7. *H contains a graph B_3 .*

Proof. By assumption H contains a copy C of C_5 . Since H has maximum degree 3, some vertex of C has a third edge. Since every edge of H is in a C_5 , and H has maximum degree three, this third edge is contained in another copy C' of C_5 which intersects C in at least an edge. Using the fact that H has girth 5, and property P6, one can show that C and C' intersect in either an edge (i.e., a 2-path), or a 3-path. That is, H contains one of the two possible broken 3-ribbons B_3 . \square

Let n be the maximum integer for which H contains a broken n -ribbon B_n , and let B a subgraph of H consisting of a copy of B_n and all of its bridges. Furthermore, assume that this B was chosen to maximise the number of bridges. Observe that B is C_5 -colourable. Because of this and the assumption that H is a core, H must have some edge not in B . This edge must be in some new C_5 , which we call C . Since H has maximum degree 3, C must intersect B in some union of paths each of whose endpoints have degree 2 in B . The following is a list of the ways in which C may intersect B .

FIGURE 4. Redrawing of $B \cup C$ in Case iii).

- (i) One edge (i.e., a 2-path.)
- (ii) Two edges.
- (iii) One 3-path.
- (iv) An edge and a 3-path.
- (v) One 4-path.
- (vi) One 5-path.

For each case listed above, we will show that one of the following must be true.

- $B \cup C$ is a (twisted) subdivided n -ribbon in which C is the link L_n .
- One of the properties P1 - P6 is contradicted.
- $B \cup C$ is a broken $(n + 1)$ -ribbon.
- $B \cup C$ is a broken n -ribbon with more bridges than B .

The lemma follows from any of these conclusions. We now go to the cases.

Case (i) The only places in B where we can have two adjacent degree two vertices, is in L_1 or L_{n-1} . If C intersects B in either one of these places, then $B \cup C$ is a broken $(n + 1)$ -ribbon.


Case (ii) One of the edges must be in L_1 and one must be in L_{n-1} . The rest of C is an edge and a 3-path between L_1 and L_{n-1} . Thus $B \cup C$ is a (twisted) subdivided n -ribbon.

Case (iii) There are only two places in B where we find two degree two vertices distance two apart.

The first place is the vertices c_i and c_{i+1} when both L_i and L_{i+1} are of type TOP or BOT. If C intersects B in the 3-path between these vertices, then the path's third vertex, a_{i+1} or b_{i+1} , contradicts property P6.

The second place is in L_1 or L_n . Assume, without loss of generality, that C intersects L_1 in a 3-path. We may assume that L_1 is of type LEFT or RIGHT. If L_1 is of type LEFT, then as in Case (ii) we have a broken $(n + 1)$ -ribbon. If L_1 is of type RIGHT, then, without loss of generality, C is the cycle $a_1 a_2 c_1 y x$, with new vertices x and y . Figure 4 gives a redrawing of $B \cup C$ in this situation which shows that $B \cup C$ is a broken n -ribbon with one more bridge than B has.

Case (iv) As in Case (iii) we immediately eliminate the possibility that the 3-path has endpoints c_i and c_{i+1} . We may assume then, that the edge is in L_1 and the 3-path is in L_{n-1} . Thus $B \cup C$ is either a (twisted) subdivided n -ribbon, or C is, up to symmetry, one of $a_1 c_1 a_n a_{n-1} c_{n-1}$ and $a_1 c_1 c_{n-1} a_{n-1} a_n$. In these last two cases, $B \cup C$ contains S_n and S_n^t respectively. (See Figure 5 for an example.)

FIGURE 5. Redrawing $B \cup C$.

Case (v) If C intersects B in a 4-path that contains three consecutive vertices of a path, then property P6 is contradicted. Avoiding this case, there are only two kinds of 4-paths in B between vertices of degree two.

The first kind is where the 4-path is between codicils of consecutive links, where one is of type TOP and one is of type BOT. If C intersects B in such a path, then $B \cup C$ is B plus another bridge.

The second kind is where, up to symmetry, the 4-path is $b_1 a_1 a_2 c_1$, where L_1 is of type RIGHT. In this case, the rest of C is $c_1 x b_1$ for some new vertex x , but then $B \cup C$ contains the 4-cycle $b_1 b_2 c_1 x b_1$. This contradicts the girth of H .

Case (vi) Again, avoiding obvious contradictions of property P6, the only possibilities contradict the girth property.

We have thus shown that H must contain S_n or S_n^t , and the only other edges H can have are bridges. □

This completes the proof of Lemma 4.6.

5. PROOF OF THEOREM 1.2

Let H be any graph of maximum degree 3 and girth 5. To prove Theorem 1.2 we must show that planar H -COL is NP -complete. By the results of Section 3, we may assume that H satisfies the properties P1 - P6. Thus H has the structure described in Lemma 4.6.

In this section we will show that planar H -COL is NP -complete. This will prove the theorem.

Lemma 4.6 gives us a structural description of H . However, not all graphs fitting this description are cores. In fact, many graphs fitting this description are C_5 -colourable. For those that are cores, we will use an indicator-type construction to show that planar H -COL is NP -complete. When H is planar, the construction is fairly straightforward. When H is not planar, an important part of the construction will be finding a planar subgraph $P(H)$ of H whose possible H -colourings are well understood. We will be able to do this because, in a manner of speaking, the description of H given by Lemma 4.6 is ‘almost’ planar.

We begin with the following lemma.

Lemma 5.1. *Let H , and its subgraph S , be as in Lemma 4.6, and assume that H is not C_5 -colourable. Let c be any vertex in S . Then there exists a planar subgraph*

$P(H)$ of S , containing c , such that any H -colouring of $P(H)$ is an injection of $P(H)$ to S .

Proof. Assume that H is not C_5 -colourable, and let c be a vertex of the subgraph S of H . Observe that S is not C_5 -colourable. Indeed, H is just the union of S with some of its bridges, so a C_5 -colouring of S would clearly imply a C_5 -colouring of H . We continue the proof in two cases: S is or isn't twisted.

In the case that S is a copy of S_n , let $P(H) = S$. We must show that the only H -colourings of S are automorphisms of S .

Let ϕ be an H -colouring of S . For any $i = 1, \dots, n$, it is easy to see that H is C_5 -colourable if we remove at least one edge from each of $C_A \cap L_i$ and $C_B \cap L_i$. Since S is not C_5 -colourable, at least one of $C_A \cap L_i$ and $C_B \cap L_i$ is in $\phi(S)$. In either case, $\phi(S)$ contains one of the edges of L_i which is in only one copy of C_5 , specifically L_i , in H . Let $\phi(e)$ be this edge. Since e is in a C_5 in S , L_i must be in $\phi(S)$. This is for all i , so S is a subgraph of $\phi(S)$. The lemma follows in this case.

In the case that S is a copy of S_n^t , by reindexing, we may assume that c is not in the n^{th} link. Since S is not C_5 colourable, and $S - (L_n \cap C_A) - (L_n \cap C_B)$ is uniquely C_5 -colourable, at least one of the planar graphs $S - (L_n \cap C_A)$ and $S - (L_n \cap C_B)$ is not C_5 -colourable. Assume, without loss of generality, that $S - (L_n \cap C_A)$ is not C_5 -colourable, and let $P(H) = S - (L_n \cap C_A)$.

By arguments similar to those in the previous case, we can show for any H -colouring ϕ of S , that all links of S except one are in $\phi(S)$, and in the last link, L_i , $\phi(L_n \cap C_A) = L_i \cap C_A$ or $L_i \cap C_B$. Again, the lemma follows. \square

Now for every plane graph G , we will construct a graph $*G$ such that

$$*G \rightarrow H \iff G \rightarrow K_3.$$

There are several cases to consider. Each case will use the following construction, and will differ only in how we choose the vertex c . In the construction we will use many copies of the cycle C_7 . Let a be a vertex of C_7 , and x and y be the two vertices that are distance 2 from a .

Construction 5.2. *Given a plane graph G , and a vertex c of the planar subgraph $P(H)$ of H , construct the planar graph $*G$ as follows.*

- Replace every edge uv of H with a copy C_{uv} of C_7 , by removing the edge uv , and identifying u and v with the copies of x and y in C_{uv} .
- For every edge uv of H attach a copy of P_{uv} of $P(H)$ to C_{uv} by identifying the copy of c in P_{uv} with the copy of a in C_{uv} .

Since G and $P(H)$ are planar, this can be done so that the resulting graph $*G$ is planar. (See Figure 6 for an example of this construction when $G = K_3$.)

Now we consider the following cases:

- (i) H has a link of type LEFT.
- (ii) All links of H are of type TOP or all are of type BOT.
- (iii) The links of H alternate between type TOP and BOT.
- (iv) The links of H alternate between two of type TOP and two of type BOT.
- (v) None of the above hold.

The proof of the backwards implication, $*G \rightarrow H \Leftarrow G \rightarrow K_3$, is very similar in each case, thus we prove it only in the first case.




FIGURE 6. Construction 5.2 for $G = K_3$.




FIGURE 7. Case (i) Mapping $*G$ to H .

Case (i) Assume that H has a link L_i of type LEFT. Let c be the codicil c_i in the planar subcore $P = P(H)$ of H . Given a planar graph G , let $*G$ be the graph returned by Construction 5.2 for this choice of P and c . We now show that

$$*G \rightarrow H \iff G \rightarrow K_3.$$

Observe that in a subdivided n -ribbon with bridges, the only vertices that are incident to two edges each of whose faces are C_5 , are codicils of links of type LEFT or RIGHT. Thus by Lemma 5.1, c must map to one of these codicils under any H -colouring of $P(H)$.

Assume there is an H -colouring ϕ of $*G$. For every uv in G , the copy of a in the subgraph C_{uv} of $*G$ is identified with the copy of c in P_{uv} , so $\phi(a)$ must be the codicil of some link of type LEFT or RIGHT. The vertices u and v , which are identified with the copies of x and y in C_{uv} , must then map to the endpoints of one of the thick edges (not necessarily in H) shown in Figure 7. (We get the same picture on a twisted link.)

The map ϕ thus induces a map ϕ' of G to the graph H^* which is the union of such ‘bowties’ over all type LEFT links. The graph H^* clearly maps K_3 , thus so does G .

On the other hand, assume that we have a homomorphism $\phi' : G \rightarrow K_3$. Define $\phi : *G \rightarrow H$ as follows. For every edge e in G , let ϕ take the copy of a in C_e to c_j . For every vertex v of G let $\phi(v)$ be c_j , a_{j+1} , or b_{j+1} if $\phi'(v)$ is 1, 2, or 3 respectively. This map can be extended to a homomorphism of $*G$ to H .




FIGURE 8. H^* for Case (ii) .




FIGURE 9. H^* for Case (iii) .

Case (ii) We may assume that all links of H are of type TOP. Let c be any codicil in $P(H)$. Given a planar graph G , let $*G$ be the graph returned by Construction 5.2. An H -colouring of $*G$ induces an H^* -colouring of G , where H^* is the graph indicated by the bold edges in Figure 8. This graph clearly maps to K_3 .

Case (iii) Let c be any codicil in $P(H)$. Given a planar graph G , let $*G$ be the graph returned by Construction 5.2. An H -colouring of $*G$ induces an H^* -colouring of G , where H^* is (a subgraph of) the graph indicated by the bold edges in Figure 9. (The broken bold edges will only be in H^* if H contains bridges.) The figure indicates a K_3 -colouring of this graph, whether or not the broken edges are present.


Case (iv) One can check that if the links of H alternate between two of type TOP and two of type BOT, then H is C_5 -colourable.

Case (v) Recall that the orbit of a vertex v in H is the set $\{u \in V(H) \mid \sigma(v) = u, \sigma \in \text{Aut}(H)\}$. Since the previous four cases cover all cases in which all of the codicils are in the same orbit, we may assume that there are at least two orbits. Furthermore, by eliminating cases (i) and (iv), we may assume that there is a link L_j of type TOP, for which both of the links L_{j-1} and L_{j+1} are of type TOP or BOT. Let c be some codicil not in the same orbit as c_j . Given a planar graph G , let $*G$ be the graph returned by Construction 5.2.

An H -colouring of $*G$ induces an H^* -colouring of G , where H^* is some graph on the vertices of H , which when restricted to the vertices of two consecutive links is some subgraph of the graphs given in Figures 7, 8, and 9.

From cases (i), (ii), and (iii), it is clear that we can always 3-colour the subgraph of H^* induced by $H - L_j$. The thick edges shown in Figure 10 are the only possible edges of the subgraph of H^* that are induced by the vertices of the links L_{j-1} , L_j and L_{j+1} . It follows from this figure that a 3-colouring of the part of H^* induced by the vertices $V(H^*) - V(L_j)$, can be extended to a 3-colouring of H^* .

We have shown that for any graph H meeting the description in Lemma 4.6, either H is C_5 -colourable, or Construction 5.2 gives us, for every planar graph G ,

FIGURE 10. H^* for Case (v) .

a planar graph *G such that

$${}^*G \rightarrow H \iff G \rightarrow K_3.$$

Since planar K_3 -COL is *NP*-complete, so is planar H -COL. This completes the proof of Theorem 1.2.

REFERENCES

- [1] J.Bondy, U.Murty, “Graph theory with applications.” American Elsevier Publishing Co., Inc., New York, 1976.
- [2] R.Brooks, *On colouring of nodes of a network*, Proc Cambridge Phil. Soc. 37 (1941) 194-197.
- [3] T.Feder, P.Hell, B.Mohar, *Acyclic homomorphisms and circular colorings of digraphs*. SIAM J. Discrete Math. 17 (2003), no. 1, 161-169.
- [4] A.Galluccio, P.Hell, J.Nešetřil, *The complexity of H -colouring of bounded degree graphs*. Discrete Math. 222 (2000), no. 1-3, 101-109.
- [5] M.Garey, D.Johnson, *Computers and Intractability* Freeman, New York, 1979.
- [6] R.Häggkvist, P.Hell. *Universality of A -mote graphs*. European J. Combin. 14 (1993) 23-27.
- [7] P.J.Heawood, *On the four-color map theorem*, Quarterly J. Pure & Applied Math 29 (1898) 270-285.
- [8] P.Hell, R.Naserasr, C.Tardif, *Complexity of Homomorphisms to Planar Graphs* Manuscript (2006).
- [9] P.Hell, J.Nešetřil, “Graphs and Homomorphisms”, Oxford University Press, 2004
- [10] P.Hell, J.Nešetřil, *On the complexity of H -colouring* J. Combin. Theory B 48 (1990) 92-100.
- [11] J.Kratochvíl, Z.Tuza, *Algorithmic complexity of list colorings*. Discrete Appl. Math. 50 (1994), no. 3, 297-302.
- [12] R.Naserasr, *Homomorphisms and Bounds*, PhD thesis, Simon Fraser University, (2003).
- [13] J.Nešetřil, P.Ossona de Mendez, *Tree-depth, subgraph colorings and homomorphism bounds*, European J. Combin. 27 (2006), 1022-1041

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF VICTORIA, PO BOX 3045
 STN CSC, VICTORIA, B.C., CANADA, V8W 3P4
E-mail address: gmacgill@math.uvic.ca, mhsiggers@gmail.com