

Real Hypersurfaces in Complex Two-Plane Grassmannians with Vanishing Lie Derivative

Young Jin Suh

Abstract. In this paper we give a characterization of real hypersurfaces of type A in a complex two-plane Grassmannian $G_2(\mathbb{C}^{m+2})$ which are tubes over totally geodesic $G_2(\mathbb{C}^{m+1})$ in $G_2(\mathbb{C}^{m+2})$ in terms of the *vanishing Lie derivative* of the shape operator A along the direction of the Reeb vector field ξ .

0 Introduction

In the geometry of real hypersurfaces there were some characterizations of homogeneous real hypersurfaces of type A_1, A_2 in complex projective space $\mathbb{C}P^m$ and of type A_0, A_1, A_2 in complex hyperbolic space $\mathbb{C}H^m$. As an example, we say that the shape operator A and the structure tensor ϕ commuting with each other, that is $A\phi = \phi A$, is a model characterization of this type hypersurface, which is a tube over a totally geodesic $\mathbb{C}P^k$ in $\mathbb{C}P^m$ (See Okumura [8]), a tube over a totally geodesic $\mathbb{C}H^k$ in $\mathbb{C}H^m$, or a horosphere in $\mathbb{C}H^m$ (See Montiel and Romero [7]).

Now let us denote by $G_2(\mathbb{C}^{m+2})$ the set of all complex two-dimensional linear subspaces in \mathbb{C}^{m+2} , which is said to be a *complex two-plane Grassmannian*. This Riemannian symmetric space $G_2(\mathbb{C}^{m+2})$ is equipped with both a Kähler structure J and a quaternionic Kähler structure \mathfrak{J} not containing J . Then for real hypersurfaces M in $G_2(\mathbb{C}^{m+2})$ we have considered the two natural geometric conditions that $[\xi] = \text{Span}\{\xi\}$ or $\mathfrak{D}^\perp = \text{Span}\{\xi_1, \xi_2, \xi_3\}$ be invariant under the shape operator A of M , where $\xi = -JN$ and $J_\nu = -J_\nu N$, $\nu = 1, 2, 3$ for a unit normal vector field N of M in $G_2(\mathbb{C}^{m+2})$. (See the details in [2, 3].)

The first result in this direction is the classification of real hypersurfaces in

$$G_2(\mathbb{C}^{m+2})$$

satisfying both conditions mentioned above. Namely, Berndt and the present author [2] have proved the following:

Theorem A *Let M be a connected real hypersurface in $G_2(\mathbb{C}^{m+2})$, $m \geq 3$. Then both $[\xi]$ and \mathfrak{D}^\perp are invariant under the shape operator of M if and only if*

- (A) *M is an open part of a tube around a totally geodesic $G_2(\mathbb{C}^{m+1})$ in $G_2(\mathbb{C}^{m+2})$, or*
- (B) *m is even, say $m = 2n$, and M is an open part of a tube around a totally geodesic $\mathbb{Q}P^n$ in $G_2(\mathbb{C}^{m+2})$.*

Received by the editors February 1, 2004; revised July 5, 2004.

This work was supported by grant Proj. No. R14-2002-003-01001-0 from the Korea Science & Engineering Foundation.

AMS subject classification: Primary: 53C40; secondary: 53C15.

©Canadian Mathematical Society 2006.

In [3], Berndt and the present author have given a characterization of real hypersurfaces of type A in Theorem A when the shape operator A of M in $G_2(\mathbb{C}^{m+2})$ commutes with the structure tensor ϕ , which is equivalent to the condition that *the Reeb flow on M is isometric*, that is $\mathcal{L}_\xi g = 0$, where \mathcal{L} (resp. g) denotes the Lie derivative (resp. the induced Riemannian metric) of M in the direction of the Reeb vector field ξ . Namely, we proved the following:

Theorem B *Let M be a connected orientable real hypersurface in $G_2(\mathbb{C}^{m+2})$, $m \geq 3$. Then the Reeb flow on M is isometric if and only if M is an open part of a tube around some totally geodesic $G_2(\mathbb{C}^{m+1})$ in $G_2(\mathbb{C}^{m+2})$.*

When the Reeb flow on M in $G_2(\mathbb{C}^{m+2})$ is *isometric* in Theorem B, we say that the Reeb vector field ξ on M is Killing. This means that the metric tensor g is invariant under the Reeb flow of ξ on M . In this paper, specifically we assert a characterization of real hypersurfaces of type A in Theorem A by another geometric Lie invariant, that is, the shape operator A of M in $G_2(\mathbb{C}^{m+2})$ is invariant under the Reeb flow on M as follows:

Main Theorem *Let M be a connected orientable real hypersurface in $G_2(\mathbb{C}^{m+2})$, $m \geq 3$. Then the Reeb flow on M satisfies $\mathcal{L}_\xi A = 0$ if and only if M is an open part of a tube around some totally geodesic $G_2(\mathbb{C}^{m+1})$ in $G_2(\mathbb{C}^{m+2})$.*

1 Riemannian Geometry of $G_2(\mathbb{C}^{m+2})$

In this section we give basic material about complex two-plane Grassmannians

$$G_2(\mathbb{C}^{m+2}),$$

for details see [2, 3]. The special unitary group $G = SU(m+2)$ acts transitively on $G_2(\mathbb{C}^{m+2})$ with stabilizer isomorphic to $K = S(U(2) \times U(m)) \subset G$. Then $G_2(\mathbb{C}^{m+2})$ can be identified with the homogeneous space G/K , which we equip with the unique analytic structure for which the natural action of G on $G_2(\mathbb{C}^{m+2})$ becomes analytic.

Now let us denote by \mathfrak{g} and \mathfrak{k} the Lie algebra of G and K , respectively, and by \mathfrak{m} the orthogonal complement of \mathfrak{k} in \mathfrak{g} with respect to the Cartan–Killing form B of \mathfrak{g} . Then $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m}$ is an $\text{Ad}(K)$ -invariant reductive decomposition of \mathfrak{g} . We put $\sigma = eK$ and identify $T_\sigma G_2(\mathbb{C}^{m+2})$ with \mathfrak{m} in the usual manner. Since B is negative definite on \mathfrak{g} , its negative restricted to $\mathfrak{m} \times \mathfrak{m}$ yields a positive definite inner product on \mathfrak{m} . By $\text{Ad}(K)$ -invariance of B this inner product can be extended to a G -invariant Riemannian metric g on $G_2(\mathbb{C}^{m+2})$.

The Lie algebra \mathfrak{k} has the direct sum decomposition $\mathfrak{k} = \mathfrak{su}(m) \oplus \mathfrak{su}(2) \oplus \mathfrak{R}$, where \mathfrak{R} is the center of \mathfrak{k} . Viewing \mathfrak{k} as the holonomy algebra of $G_2(\mathbb{C}^{m+2})$, the center \mathfrak{R} induces a Kähler structure J and the $\mathfrak{su}(2)$ -part a quaternionic Kähler structure \mathfrak{J} on $G_2(\mathbb{C}^{m+2})$. If J_1 is any almost Hermitian structure in \mathfrak{J} , then $JJ_1 = J_1J$, and JJ_1 is a symmetric endomorphism with $(JJ_1)^2 = I$ and $\text{tr}(JJ_1) = 0$.

A canonical local basis J_1, J_2, J_3 of \mathfrak{J} consists of three local almost Hermitian structures J_ν in \mathfrak{J} such that $J_\nu J_{\nu+1} = J_{\nu+2} = -J_{\nu+1} J_\nu$, where the index is taken modulo three. Since \mathfrak{J} is parallel with respect to the Riemannian connection $\bar{\nabla}$ of

$(G_2(\mathbb{C}^{m+2}), g)$, there exist for any canonical local basis J_1, J_2, J_3 of \mathfrak{J} three local one-forms q_1, q_2, q_3 such that

$$(1.1) \quad \bar{\nabla}_X J_\nu = q_{\nu+2}(X) J_{\nu+1} - q_{\nu+1}(X) J_{\nu+2}$$

for all vector fields X on $G_2(\mathbb{C}^{m+2})$.

2 Some Fundamental Formulas for Real Hypersurfaces in $G_2(\mathbb{C}^{m+2})$

In this section we derive some basic formulae from the Codazzi equation for a real hypersurface M in $G_2(\mathbb{C}^{m+2})$.

The induced Riemannian metric on M will also be denoted by g , and ∇ denotes the Riemannian connection of (M, g) . The Kähler structure J of $G_2(\mathbb{C}^{m+2})$ induces an almost contact metric structure (ϕ, ξ, η, g) on M . Furthermore, let J_1, J_2, J_3 be a canonical local basis of \mathfrak{J} . Then each J_ν induces an almost contact metric 3-structure $(\phi_\nu, \xi_\nu, \eta_\nu, g)$ on M . Using the expression for the curvature tensor \bar{R} of $G_2(\mathbb{C}^{m+2})$ given in [2] and [3], the Codazzi equation becomes

$$\begin{aligned} (\nabla_X A)Y - (\nabla_Y A)X &= \eta(X)\phi Y - \eta(Y)\phi X - 2g(\phi X, Y)\xi \\ &+ \sum_{\nu=1}^3 \{ \eta_\nu(X)\phi_\nu Y - \eta_\nu(Y)\phi_\nu X - 2g(\phi_\nu X, Y)\xi_\nu \} \\ &+ \sum_{\nu=1}^3 \{ \eta_\nu(\phi X)\phi_\nu \phi Y - \eta_\nu(\phi Y)\phi_\nu \phi X \} \\ &+ \sum_{\nu=1}^3 \{ \eta(X)\eta_\nu(\phi Y) - \eta(Y)\eta_\nu(\phi X) \} \xi_\nu. \end{aligned}$$

The following identities can be proved in a straightforward method and will be used frequently in subsequent calculations, (see [10, 11]):

$$\begin{aligned} (2.1) \quad \phi_{\nu+1}\xi_\nu &= -\xi_{\nu+2}, \quad \phi_\nu\xi_{\nu+1} = \xi_{\nu+2}, \\ \phi\xi_\nu &= \phi_\nu\xi, \quad \eta_\nu(\phi X) = \eta(\phi_\nu X), \\ \phi_\nu\phi_{\nu+1}X &= \phi_{\nu+2}X + \eta_{\nu+1}(X)\xi_\nu, \\ \phi_{\nu+1}\phi_\nu X &= -\phi_{\nu+2}X + \eta_\nu(X)\xi_{\nu+1}, \end{aligned}$$

where the index ν denotes $\nu = 1, 2, 3$.

Now let us put

$$JX = \phi X + \eta(X)N, \quad J_\nu X = \phi_\nu X + \eta_\nu(X)N$$

for any tangent vector X of a real hypersurface M in $G_2(\mathbb{C}^{m+2})$, where N denotes a unit normal vector field of M in $G_2(\mathbb{C}^{m+2})$. Then from this and the formulas (1.1)

and (2.1) we have that

$$(2.2) \quad (\nabla_X \phi)Y = \eta(Y)AX - g(AX, Y)\xi, \quad \nabla_X \xi = \phi AX,$$

$$(2.3) \quad \nabla_X \xi_\nu = q_{\nu+2}(X)\xi_{\nu+1} - q_{\nu+1}(X)\xi_{\nu+2} + \phi_\nu AX,$$

$$(2.4) \quad (\nabla_X \phi_\nu)Y = -q_{\nu+1}(X)\phi_{\nu+2}Y + q_{\nu+2}(X)\phi_{\nu+1}Y + \eta_\nu(Y)AX - g(AX, Y)\xi_\nu.$$

Moreover, from $JJ_\nu = J_\nu J$, $\nu = 1, 2, 3$, it follows that

$$(2.5) \quad \phi\phi_\nu X = \phi_\nu\phi X + \eta_\nu(X)\xi - \eta(X)\xi_\nu.$$

3 Proof of the Main Theorem

Before giving the proof of our theorem, let us determine which of the model hypersurfaces given in Theorem A satisfy the formula $\mathcal{L}_\xi A = 0$. First note that

$$\begin{aligned} (\mathcal{L}_\xi A)X &= \mathcal{L}_\xi(AX) - A\mathcal{L}_\xi X \\ &= \nabla_\xi(AX) - \nabla_{AX}\xi - A(\nabla_\xi X - \nabla_X \xi) \\ &= (\nabla_\xi A)X - \nabla_{AX}\xi + A\nabla_X \xi \\ &= (\nabla_\xi A)X - \phi A^2 X + A\phi AX \\ &= 0 \end{aligned}$$

for any vector field X on M . Then the assumption $\mathcal{L}_\xi A = 0$ holds if and only if $(\nabla_\xi A)X = \phi A^2 X - A\phi AX$. In this section we will show that only a tube over a totally geodesic $G_2(\mathbb{C}^{m+1})$ in $G_2(\mathbb{C}^{m+2})$ satisfies the formula $\mathcal{L}_\xi A = 0$.

Now let us consider a real hypersurfaces of type A, that is, a tube over a totally geodesic $G_2(\mathbb{C}^{m+1})$ in $G_2(\mathbb{C}^{m+2})$. By Proposition A of [11] we know that $\xi \in \mathfrak{D}^\perp$ and that the shape operator A and the structure tensor ϕ commute with each other. Then this implies that ξ is principal, that is, $A\xi = \alpha\xi$, where $\alpha = \sqrt{8} \cot(\sqrt{8}r)$. Differentiating this one, by (2.2) we have

$$\begin{aligned} (\nabla_X A)\xi &= \nabla_X(A\xi) - A\nabla_X \xi \\ &= \alpha\nabla_X \xi - A\nabla_X \xi \\ &= \alpha\phi AX - A\phi AX. \end{aligned}$$

On the other hand, by the equation of Codazzi and the assumption of $\mathcal{L}_\xi A = 0$ we have

$$\begin{aligned} (3.1) \quad (\nabla_X A)\xi &= -\phi X - \sum_{\nu} \{\eta_\nu(\xi)\phi_\nu X - \eta_\nu(X)\phi_\nu \xi\} \\ &\quad - 3 \sum_{\nu} \eta_\nu(\phi X)\xi_\nu + \phi A^2 X - A\phi AX, \end{aligned}$$

where \sum_ν denotes the summation from $\nu = 1$ to 3. Then from these two formulas we have

$$(3.2) \quad \begin{aligned} \alpha\phi AX - A\phi AX &= -\phi X - \phi_1 X + \sum_\nu \eta_\nu(X)\phi_\nu \xi \\ &\quad - 3 \sum_\nu \eta_\nu(\phi X)\xi_\nu + \phi A^2 X - A\phi AX. \end{aligned}$$

Now let us check case by case whether two sides in (3.2) are equal to each other as follows:

Case 1. $X = \xi = \xi_1$

In this case it can be easily checked that two sides are equal to each other.

Case 2. $X = \xi_2, \xi_3$

Then we may put $A\xi_2 = \beta\xi_2$, $A\xi_3 = \beta\xi_3$, where $\beta = \sqrt{2} \cot(\sqrt{2}r)$. Then by putting $X = \xi_2$ in (3.2) we have

$$\alpha\beta\phi\xi_2 = -3\xi_3 - 3 \sum_\nu \eta_\nu(\phi\xi_2)\xi_\nu + \beta^2\phi\xi_2.$$

From this we know $\beta(\alpha - \beta)\phi\xi_2 = 2\xi_3$, which implies that both sides are equal to $2\xi_3$.

Case 3. $X \in T_\lambda = \{X | X \perp \mathbb{H}\xi, \phi X = \phi_1 X\}$

Then by putting $X \in T_\lambda$, $\lambda = -\sqrt{2} \tan(\sqrt{2}r)$ in (3.2) we have

$$\alpha\lambda\phi X = -\phi X - \phi_1 X + \sum_\nu \eta_\nu(X)\phi_\nu \xi - 3 \sum_\nu \eta_\nu(\phi X)\xi_\nu + \lambda^2\phi X,$$

from this it implies that $\lambda(\alpha - \lambda)\phi X = -2\phi X$. This gives our assertion.

Case 4. $X \in T_\mu = \{X | X \perp \mathbb{H}\xi, \phi X = -\phi_1 X\}$

By putting $X \in T_\mu$, $\mu = 0$, in (3.2) we know that both sides are all vanishing.

Accordingly we conclude that real hypersurfaces of type A in Theorem A satisfy $\mathcal{L}_\xi A = 0$. In this section we are going to give the complete classification of real hypersurfaces M in $G_2(\mathbb{C}^{m+2})$ satisfying $\mathcal{L}_\xi A = 0$.

Now let us take an orthonormal basis $\{e_1, \dots, e_{4m-1}\}$ for the tangent space $T_x M$, $x \in M$, of a real hypersurface M in $G_2(\mathbb{C}^{m+2})$. Then by the equation of Codazzi we may put

$$\begin{aligned}
 (3.3) \quad (\nabla_{e_i} A)X - (\nabla_X A)e_i &= \eta(e_i)\phi X - \eta(X)\phi e_i - 2g(\phi e_i, X)\xi \\
 &+ \sum_{\nu=1}^3 \{ \eta_\nu(e_i)\phi_\nu X - \eta_\nu(X)\phi_\nu e_i - 2g(\phi_\nu e_i, X)\xi_\nu \} \\
 &+ \sum_{\nu=1}^3 \{ \eta_\nu(\phi e_i)\phi_\nu \phi X - \eta_\nu(\phi X)\phi_\nu \phi e_i \} \\
 &+ \sum_{\nu=1}^3 \{ \eta(e_i)\eta_\nu(\phi X) - \eta(X)\eta_\nu(\phi e_i) \} \xi_\nu,
 \end{aligned}$$

from which, together with the formulas (2.1) and (2.5) it follows that

$$\begin{aligned}
 (3.4) \quad \sum_{i=1}^{4m-1} g((\nabla_{e_i} A)X, \phi e_i) \\
 &= -(4m-2)\eta(X) \\
 &+ \sum_{\nu} \{ g(\phi_\nu X, \phi \xi_\nu) + \eta_\nu(X) \operatorname{Tr} \phi \phi_\nu + 2g(\phi_\nu^2 \xi, X) \} \\
 &- \sum_{\nu} g(\phi_\nu \phi X, \phi \phi_\nu \xi) - \sum_{\nu} \eta(X) g(\phi \xi_\nu, \phi \xi_\nu) \\
 &= -(4m-2)\eta(X) - 3\eta(X) + \sum_{\nu} \eta_\nu(\xi)\eta_\nu(X) + \sum_{\nu} \eta_\nu(X) \operatorname{Tr} \phi \phi_\nu \\
 &- \sum_{\nu} \eta_\nu(\xi) g(\phi_\nu \phi X, \xi) - 3\eta(X) + \sum_{\nu} \eta^2(\xi_\nu)\eta(X) \\
 &= -4(m+1)\eta(X) + 2 \sum_{\nu} \eta_\nu(\xi)\eta_\nu(X) + \sum_{\nu} \eta_\nu(X) \operatorname{Tr} \phi \phi_\nu,
 \end{aligned}$$

where in the second equality we have used the formulas

$$\begin{aligned}
 \sum_{\nu} g(\phi_\nu \phi X, \phi \phi_\nu \xi) &= \sum_{\nu} \eta_\nu(\xi) g(\phi_\nu \phi X, \xi), \\
 \sum_{\nu} g(\phi \xi_\nu, \phi \xi_\nu) &= 3\eta(X) - \sum_{\nu} \eta^2(\xi_\nu)\eta(X), \\
 \sum_{\nu} g(\phi_\nu^2 \xi, X) &= -3\eta(X) + \sum_{\nu} \eta_\nu(\xi)\eta_\nu(X),
 \end{aligned}$$

and respectively in the third equality,

$$- \sum_{\nu} \eta_\nu(\xi) g(\phi_\nu \phi X, \xi) = \sum_{\nu} \eta_\nu(\xi)\eta_\nu(X) - \sum_{\nu} \eta(X)\eta_\nu^2(\xi).$$

Now let us denote by U the vector $\nabla_\xi \xi = \phi A \xi$. Then by (2.2) its derivative can be given by

$$\nabla_{e_i} U = \eta(A \xi) A e_i - g(A e_i, A \xi) \xi + \phi(\nabla_{e_i} A) \xi + \phi A \nabla_{e_i} \xi.$$

Then its divergence is given by

$$(3.5) \quad \begin{aligned} \operatorname{div} U &= \sum_i g(\nabla_{e_i} U, e_i) \\ &= h\eta(A\xi) - \eta(A^2\xi) - g((\nabla_{e_i} A)\xi, \phi e_i) - g(\phi A e_i, A\phi e_i), \end{aligned}$$

where h denotes the trace of the shape operator of M in $G_2(\mathbb{C}^{m+2})$.

Now we calculate the squared norm of the following:

$$(3.6) \quad \begin{aligned} \|\phi A - A\phi\|^2 &= \sum_i g((\phi A - A\phi)e_i, (\phi A - A\phi)e_i) \\ &= \sum_{i,j} g((\phi A - A\phi)e_i, e_j) g((\phi A - A\phi)e_i, e_j) \\ &= \sum_{i,j} \{g(\phi A e_j, e_i) + g(\phi A e_i, e_j)\} \{g(\phi A e_j, e_i) + g(\phi A e_i, e_j)\} \\ &= 2 \sum_{i,j} g(\phi A e_j, e_i) g(\phi A e_j, e_i) + 2 \sum_{i,j} g(\phi A e_j, e_i) g(\phi A e_i, e_j) \\ &= -2 \sum_j g(\phi A e_j, A\phi e_j) + 2 \sum_j g(\phi A e_j, \phi A e_j) \\ &= 2 \operatorname{Tr} A^2 - 2h\eta(A\xi) + 2 \sum_i g((\nabla_{e_i} A)\xi, \phi e_i) + 2 \operatorname{div} U, \end{aligned}$$

where \sum_i (resp. $\sum_{i,j}$) denotes the summation from $i = 1$ to $i = 4m - 1$ (resp. from $i, j = 1$ to $4m - 1$) and in the final equality we have used (3.5). From this together with the formula (3.4) it follows that

$$(3.7) \quad \begin{aligned} \operatorname{div} U &= \frac{1}{2} \|\phi A - A\phi\|^2 - \operatorname{Tr} A^2 + \alpha h \\ &\quad + 4(m+1) - 2 \sum_{\nu} \eta_{\nu}^2(\xi) - \sum_{\nu} \eta_{\nu}(\xi) \operatorname{Tr} \phi \phi_{\nu}. \end{aligned}$$

From this formula, together with the assumption $\mathcal{L}_{\xi} A = 0$ we want to show that the structure tensor ϕ and the shape operator A commute with each other, that is, $\phi A - A\phi = 0$. Then by Theorem B we are able to assert that M is a tube over a totally geodesic $G_2(\mathbb{C}^{m+1})$ in $G_2(\mathbb{C}^{m+2})$.

Now let us take an inner product (3.1) with the Reeb vector field ξ . Then we have

$$(3.8) \quad \begin{aligned} g((\nabla_X A)\xi, \xi) &= - \sum_{\nu} \eta_{\nu}(\xi) \eta_{\nu}(\phi X) - 3 \sum_{\nu} \eta_{\nu}(\phi X) \eta_{\nu}(\xi) - g(A\phi A X, \xi) \\ &= -4 \sum_{\nu} \eta_{\nu}(\xi) \eta_{\nu}(\phi X) + g(AX, U). \end{aligned}$$

On the other hand, by the almost contact structure ϕ we have

$$\phi U = \phi^2 A\xi = -A\xi + \eta(A\xi)\xi = -A\xi + \alpha\xi,$$

where the function α denotes $\eta(A\xi)$. From this, differentiating and using (2.2) gives

$$(3.9) \quad (\nabla_X \phi)U + \phi \nabla_X U = -(\nabla_X A)\xi - A \nabla_X \xi + (X\alpha)\xi + \alpha \nabla_X \xi \\ - g(AX, U)\xi + \phi \nabla_X U.$$

Then by taking an inner product (3.9) with ξ and using $U = \phi A\xi$ we have

$$g((\nabla_X A)\xi, \xi) = g(AX, U) + X\alpha - g(\nabla_X \xi, A\xi) \\ = 2g(AX, U) + X\alpha.$$

From this, together with (3.8), we have

$$(3.10) \quad g(AX, U) + 4 \sum_{\nu} \eta_{\nu}(\xi) \eta_{\nu}(\phi X) + X\alpha = 0.$$

Now substituting (3.1) and (3.10) into (3.9) and using (2.2), we have

$$(3.11) \quad 4 \sum_{\nu} \eta_{\nu}(\xi) \eta_{\nu}(\phi X) + \phi \nabla_X U = \phi X + \sum_{\nu} \{ \eta_{\nu}(\xi) \phi_{\nu} X - \eta_{\nu}(X) \phi_{\nu} \xi \} \\ + 3 \sum_{\nu} \eta_{\nu}(\phi X) \xi_{\nu} - \phi A^2 X + \alpha \phi AX.$$

Then the above equation can be rewritten as follows:

$$\phi \nabla_X U = \phi X - \phi A^2 X + \alpha \phi AX + \sum_{\nu} \{ \eta_{\nu}(\xi) \phi_{\nu} X - \eta_{\nu}(X) \phi_{\nu} \xi \} \\ + 3 \sum_{\nu} \eta_{\nu}(\phi X) \xi_{\nu} - 4 \sum_{\nu} \eta_{\nu}(\xi) \eta_{\nu}(\phi X) \xi.$$

From this, summing up from 1 to $4m - 1$ for an orthonormal basis of $T_x M$, $x \in M$, we have

$$(3.12) \quad \sum_i g(\phi \nabla_{e_i} U, \phi e_i) = \operatorname{div} U + \|U\|^2 \\ = (4m - 2) - \operatorname{Tr} A^2 + \eta(A^2 \xi) + \alpha \{ \operatorname{Tr} A - \alpha \} \\ - \sum_{\nu} \eta_{\nu}(\xi) \operatorname{Tr} \phi \phi_{\nu} \\ - \sum_{\nu} g(\phi_{\nu} \xi, \phi_{\nu} \xi) + 3 \sum_{\nu} g(\xi_{\nu}, \xi_{\nu}) - 3 \sum_{\nu} \eta^2(\xi_{\nu}),$$

where in the first equality we have used the notion of $\operatorname{div} U$. Then it follows that

$$\begin{aligned}
 (3.13) \quad \operatorname{div} U &= (4m - 2) - \operatorname{Tr} A^2 + \alpha \operatorname{Tr} A - \sum_{\nu} \eta_{\nu}(\xi) \operatorname{Tr} \phi \phi_{\nu} \\
 &\quad - \left\{ 3 - \sum_{\nu} \eta(\xi_{\nu}) \eta(\xi_{\nu}) \right\} + 9 - 3 \sum_{\nu} \eta^2(\xi_{\nu}) \\
 &= 4(m + 1) + \alpha h - \operatorname{Tr} A^2 - \sum_{\nu} \eta_{\nu}(\xi) \operatorname{Tr} \phi \phi_{\nu} - 2 \sum_{\nu} \eta^2(\xi_{\nu}),
 \end{aligned}$$

where we have used $\|U\|^2 = \|A\xi\|^2 - \alpha^2$ in (3.12).

Now if we compare (3.7) with the formula (3.13), we finally assert that the squared norm $\|A\phi - \phi A\|^2$ vanishes, that is, the structure tensor ϕ and the shape operator A commute with each other. Then by Theorem B in the introduction we are able to assert that M is a tube over a totally geodesic $G_2(\mathbb{C}^{m+1})$ in $G_2(\mathbb{C}^{m+2})$. This completes the proof of our Main Theorem. ■

Remark 3.1 Let M be a real hypersurface in complex two-plane Grassmannians $G_2(\mathbb{C}^{m+2})$ satisfying $\mathcal{L}_{\xi}\phi = 0$. Then it is not difficult to show that the conditions $\mathcal{L}_{\xi}\phi = 0$ and $\mathcal{L}_{\xi}A = 0$ are equivalent. So we remark here that a real hypersurface M in $G_2(\mathbb{C}^{m+2})$ satisfying $\mathcal{L}_{\xi}\phi = 0$ is also congruent to a tube over a totally geodesic $G_2(\mathbb{C}^{m+1})$ in $G_2(\mathbb{C}^{m+2})$.

Remark 3.2 In paper [10] due to the present author we have proved some non-existence properties for real hypersurfaces in $G_2(\mathbb{C}^{m+2})$ with parallel shape operator $\nabla A = 0$. Also in [11] we have investigated some real hypersurfaces M in $G_2(\mathbb{C}^{m+2})$ when the structure tensors ϕ_{ν} , $\nu = 1, 2, 3$, commute with the shape operator A of M in $G_2(\mathbb{C}^{m+2})$.

Acknowledgement The author would like to express his sincere gratitude to the referee for careful reading of the manuscript and valuable comments.

References

- [1] J. Berndt, *Real hypersurfaces in quaternionic space forms*. J. Reine Angew. Math. **419**(1991), 9–26.
- [2] J. Berndt and Y. J. Suh, *Real hypersurfaces in complex two-plane Grassmannians*. Monat. Math. **127**(1999), 1–14.
- [3] ———, *Isometric flows on real hypersurfaces in complex two-plane Grassmannians*. Monat. Math. **137**(2002), 87–98.
- [4] T. E. Cecil and P. J. Ryan, *Focal sets and real hypersurfaces in complex projective space*. Trans. Amer. Math. Soc. **269**(1982), 481–499.
- [5] M. Kimura, *Real hypersurfaces and complex submanifolds in complex projective space*. Trans. Amer. Math. Soc. **296**(1986), 137–149.
- [6] A. Martinez and J. D. Pérez, *Real hypersurfaces in quaternionic projective space*. Ann. Mat. Pura Appl. **145**(1986), 355–384.
- [7] S. Montiel and A. Romero, *On some real hypersurfaces of a complex hyperbolic space*. Geom. Dedicata **20**(1986), 245–261.
- [8] M. Okumura, *On some real hypersurfaces of a complex projective space*. Trans. Amer. Math. Soc. **212**(1975), 355–364.

- [9] J. D. Pérez and Y. J. Suh, *Real hypersurfaces of quaternionic projective space satisfying $\nabla_{U_i} R = 0$* . Differential Geom. and Appl. **7**(1997), 211–217.
- [10] Y. J. Suh, *Real hypersurfaces in complex two-plane Grassmannians with parallel shape operator*. Bull. Austral. Math. Soc. **67**(2003), 493–502.
- [11] ———, *Real hypersurfaces in complex two-plane Grassmannians with commuting shape operator*. Bull. Austral. Math. Soc. **68**(2003), 379–393.
- [12] K. Yano and M. Kon, *CR-submanifolds of Kaehlerian and Sasakian manifolds*. Birkhäuser, Boston, Basel, Struttgart, 1983.

Kyungpook National University
Department of Mathematics
Taegu 702-701
Korea
e-mail: yjsuh@mail.knu.ac.kr